

Global N-glycan Site Occupancy of HIV-1 gp120 by Metabolic Engineering and High-Resolution Intact Mass Spectrometry

Weston B. Struwe, Alexandra Stuckmann, Anna-Janina Behrens, Kevin Pagel, and Max Crispin

ACS Chem. Biol., **Just Accepted Manuscript** • DOI: 10.1021/acscchembio.6b00854 • Publication Date (Web): 16 Dec 2016

Downloaded from <http://pubs.acs.org> on December 22, 2016

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



Global N-glycan Site Occupancy of HIV-1 gp120 by Metabolic Engineering and High-Resolution Intact Mass Spectrometry

Weston B. Struwe*[§], Alexandra Stuckmann[†], Anna-Janina Behrens[§], Kevin Pagel[†], Max Crispin*^{§¶}

[§]Oxford Glycobiology Institute, Department of Biochemistry, University of Oxford, Oxford, OX1 3QU, United Kingdom

[†]Free University Berlin, Department of Chemistry and Biochemistry, Takustrasse 3, 14195 Berlin, Germany

[¶]Department of Immunology and Microbial Science, The Scripps Research Institute, La Jolla, CA 92037, USA

Supporting Information Placeholder

ABSTRACT: A vital step in HIV vaccine development strategies has been the observation that some infected individuals generate broadly neutralizing antibodies that target the glycans on the surface of HIV-1 gp120. These antibodies target glycan epitopes on viral envelope spikes and yet the positions and degree of occupancy of glycosylation sites is diverse. Therefore, there is a need to understand glycosylation occupancy on recombinant immunogens. The sheer number of potential glycosylation sites and degree of chemical heterogeneity impedes assessing the global sequon occupancy of gp120 glycoforms. Here, we trap the glycan processing of recombinant gp120 to generate homogenous glycoforms, facilitating occupancy assessment by intact mass spectrometry. We show that gp120 monomers of the BG505 strain contain either fully occupied sequons or missing one and sometimes two glycans across the molecule. This biosynthetic engineering approach enables the analysis of therapeutically important glycoproteins otherwise recalcitrant to analysis by native mass spectrometry.

The human immunodeficiency virus (HIV-1) viral spikes have an extensive and dense coat of N-linked glycans that act to shield the underlying protein from antibody recognition¹⁻⁷. The attachment glycoprotein (gp120) within these spikes is a key target for antibody-mediated neutralization^{8, 9}. Over time, many infected individuals produce broadly neutralizing antibodies (bnAbs) against HIV viral spike epitopes. These antibodies offer broad protection to infection in passive transfer experiments¹⁰ and eliciting bnAbs by vaccination with viral spike mimics is a key goal in the control of the pandemic¹¹. The epitopes targeted by the majority of bnAbs contain one or more glycans^{12, 13}. Although gp120 N-glycans are largely restricted to high-mannose type, the number and location of N-glycans may change during the viral life span^{1, 14}. Finally, glycan occupancy of key sites modulates the development of a broad antibody response against heterologous viruses^{15, 16}. To this end, it is important that recombinant candidate immunogens are fully characterized¹⁷, invoke a suitable T-cell response¹⁸ and efficiently display target bnAb epitopes to B-cells^{11, 19, 20}.

The extensive role of glycans in forming the epitopes of bnAbs and the emerging importance of viral site occupancy has necessitated detailed glycosylation analysis of recombinant mimics of the viral spike. This is important in guiding immunogen design and also in evaluating biotherapeutic glycoproteins for use in the clinic. Glycoproteins are known to consist of an ensemble of 'gly-

coforms'. These arise during cellular biosynthesis and the heterogeneity is driven by variable occupancy of the glycan sites and the chemical heterogeneity that arises from the action of an array of glycosidases and glycosyltransferases in the Golgi apparatus²¹. Partial occupancy of N-glycan sequons can have substantial impact on biological activity and is an important parameter in the characterization of biologics.

Significant progress has been made in site-specific analysis of gp120 glycosylation²²⁻²⁴ but little is known about the overall occupancy of glycosylation sites. While glycopeptide analysis can reveal the occupancy of any particular site^{25, 26}, measuring the overall distribution of partially occupied sites across the spectrum of glycoproteins has not been tractable by current methods. As such, glycan heterogeneity obscures global occupancy information that could be derived by intact mass spectrometry (MS)²⁷.

Here, we circumvent this barrier by using metabolic engineering with a potent α -mannosidase inhibitor, kifunensine²⁸, to homogenize the processing of N-linked glycans on recombinant gp120 (BG505 strain) transiently expressed in human embryonic kidney (HEK) 293F cells (Fig. 1A and Fig. S1).

Homogeneous gp120 glycoforms could be resolved using a




Figure 1. (A) Expression strategy to produce an oligomannose-type glycoform of gp120. Kifunensine inhibits endoplasmic reticulum (ER) and Golgi mannosidase I during recombinant HIV gp120 expression resulting in predominantly Man₉GlcNAc₂ (Man9) N-glycans. (B) ESI-MS of N-linked glycans released by protein N-glycanase F. (C) Tandem mass spectrometry of negative N-glycan ions (diagnostic ions for each isomer are in orange or purple). Green circles, mannose; Blue squares, GlcNAc.

modified high-resolution Orbitrap mass spectrometer designed to evaluate high molecular weight proteins and their complexes²⁹. High-resolution MS has been applied to glycoproteins with only one³⁰ or two glycan sites^{31, 32}, but not to highly glycosylated proteins due to overlapping glycoforms.

Kifunensine has previously been used to augment the crystallization of glycoproteins and is sufficiently potent to almost entirely eliminate chemical heterogeneity of N-linked glycosylation³³⁻³⁵. MS of released N-linked glycans from BG505 gp120 expressed in the presence of kifunensine shows a spectrum dominated by Man₉GlcNAc₂ (Man9) with only a trace of Man₈GlcNAc₂ (Man8) (Fig. 1B). Tandem MS reveals the known isomers of the mammalian glycosylation pathway (Fig. 1C)³⁶. This is consistent with




Figure 2. Intact mass spectrum of BG505 gp120 transiently expressed in the presence of kifunensine. (A) Spectrum revealing six distinct ion species within each charge state (inset). A 162 Da mass difference corresponding to Man9/Man8 microheterogeneity is present within each species (inset, green circles). The major ions are separated by an 1864 Da mass shifts (blue, inset) equal to a single Man9. (B) Positive ion MS spectrum of released O-glycans identify mono- and di-sialyl core-1 structures that correspond to the peak shifts observed in the intact spectrum (pink circles, top). Peaks marked with asterisks are non-carbohydrate contaminants. (C) Deconvoluted spectrum to quantify the major peaks in A identify fully occupied $n = 24$ or $n = 23$ glycoforms with sialyl core-1 \pm disialyl core-1 O-glycans.

efficient blockade of both endoplasmic reticulum and type-I Golgi-resident α -mannosidase activity.

Native and deconvoluted mass spectra of the resulting glycan-engineered gp120 revealed a charge state distribution spanning 15-19+ (Fig. 2A). Within each charge state, six species were observed, with an evident mass shift between the three major peaks matching the mass of a single Man9 (1864 Da) demonstrating variability in occupied glycan sites in the intact gp120 (Fig. 2A, inset). The gp120 structure is dominated by Man9 glycans, the cumulative effect of the low levels of Man8 structures gives rise to an evident hexose series within each major peak demonstrating Man8 and Man9 microheterogeneity.

Further examination of the spectra reveals mass shifts corresponding to variable modification with O-glycans (Fig. 2A, pink circles). These observations are consistent with previous glycopeptide analysis where recombinant gp120 contained O-linked glycans^{24, 26}. Notably, the measured mass of gp120 (101 kDa) was 1.6 kDa greater than that corresponding to the gp120 peptide backbone (~54.6 kDa) with 23 or 24 Man9 N-glycans (97.5 and 99.4 kDa, respectively) (Fig. S2). Analysis of released O-glycans confirmed mono- and di-sialylated core-1 structures (Fig. 2B) and accounts for the observed intact gp120 masses. Therefore, recombinant gp120 contains a single sialyl core-1 (656 Da) \pm di-sialyl core-1 (947 Da) O-glycans. From the deconvoluted spectra (Fig. 2C) we can conclude that the dominant species matched to both fully occupied and $n-1$ (i.e. lacking a single N-glycan) gp120 protein moieties with variable O-glycosylation. The approximate relative abundance of gp120 glycoforms were 40.6 % fully occupied, 41.6% ($n-1$), and 17.8% ($n-2$).

To confirm the assigned peaks to N-glycans, we measured a BG505 gp120 variant mutated at a single highly conserved glycosylation site (N332) that is a critical component of many bnAb epitopes. Expectedly, the N332A mass spectrum lacked the $n = 24$ peaks, most evident from the 16+ and 17+ charge states (Fig. 3).




Figure 3. Intact mass spectra of N332A gp120 (black) and wild-type (WT) gp120 (blue) expressed in the presence of kifunensine. The number of N-glycosylation sites at each cluster of peaks is indicated (n). WT and the N332A mutant of BG505 gp120 have a maximum N-glycosylation site occupancy potential of $n = 24$ and 23, respectively.

The depletion of two clusters of peaks per charge state also substantiates the presence both O-glycans per gp120 monomer. Interestingly, the relative occupancy was maintained with $n-1$ and $n-2$ as the major glycoforms suggesting that N332 is largely completely occupied and that the global loss of glycan equivalents arises from the cumulative partial absence of glycans across other sites (Fig. S3).

Native high-resolution mass spectrometry holds enormous promise for glycoprotein characterization, but has only been demonstrated on relatively simple systems and has failed for more

complex targets³⁰⁻³². We have chosen one of the most heavily glycosylated glycoproteins in nature which is currently under investigation for clinical use³⁷. Our results demonstrate that the assessment of global occupancy of glycosylation sites is tractable by intact mass spectrometry with biosynthetic engineering to eliminate heterogeneity arising from glycan processing. The occupancy information is ostensibly preserved as the engineered glycan processing occurs downstream of the initial oligosaccharyl-transferase that initiates N-glycosylation. Intact MS can rapidly assess the presence and structure of variable O-glycosylation which may otherwise be obscured by masses arising from heterogeneous N-link glycosylation. Detection of O-linked glycosylation remains an active area of investigation in the assessment of HIV immunogens^{24, 38}. Furthermore, this approach is widely applicable for any N-glycosylated protein and will prove valuable in biotherapeutic characterization. The clinical use of complex glycoproteins requires a detailed understanding of structure-function relationships and the monitoring of key critical quality attributes, such as glycan site occupancy³⁹. In particular, we expect assessment of global site occupancy to emerge as an important parameter in HIV immunogen production systems as these are assessed for the manufacture of clinical grade material¹⁷.

Glycan engineering has already accelerated developments in structural biology by preventing glycan heterogeneity impeding glycoprotein crystallization³³. We suggest that glycan engineering could similarly solve the 'glycosylation problem' in native mass spectrometry by simplifying the spectra for any studies dealing with heavily glycosylated glycoproteins and their interactions.

ASSOCIATED CONTENT

Supporting Information

Materials and methods, Figures S1-S3. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Authors

*weston.struwe@bioch.ox.ac.uk & max.crispin@bioch.ox.ac.uk

Notes

The subject matter of this paper is related to United Kingdom Patent Application No. 1620930.6.

ACKNOWLEDGMENTS

We thank Prof. Dame C. Robinson FRS (University of Oxford) for instrument access and Prof. I.A. Wilson FRS (The Scripps Research Institute) and Prof. R.A. Dwek FRS (University of Oxford) for useful discussions. W.B.S. is a Research Associate at University College, Oxford. A.-J.B. is a recipient of a Chris Scanlan Memorial Scholarship from Corpus Christi College, Oxford. This work was supported the International AIDS Vaccine Initiative Neutralizing Antibody Center CAVD grant (Glycan characterization and Outer Domain glycoform design) and the Scripps CHAVI-ID (1UM1A1100663).

REFERENCES



1. Scanlan, C. N., Offer, J., Zitzmann, N., and Dwek, R. A. (2007) Exploiting the defensive sugars of HIV-1 for drug and vaccine design, *Nature* **446**, 1038-1045.
2. Lyumkis, D., Julien, J. P., de Val, N., Cupo, A., Potter, C. S., Klasse, P. J., Burton, D. R., Sanders, R. W., Moore, J. P., Carragher, B., Wilson, I. A., and Ward, A. B. (2013) Cryo-EM structure of a fully glycosylated soluble cleaved HIV-1 envelope trimer, *Science* **342**, 1484-1490.
3. Julien, J. P., Cupo, A., Sok, D., Stanfield, R. L., Lyumkis, D., Deller, M. C., Klasse, P. J., Burton, D. R., Sanders, R. W., Moore, J. P., Ward, A.

B., and Wilson, I. A. (2013) Crystal structure of a soluble cleaved HIV-1 envelope trimer, *Science* **342**, 1477-1483.

4. Wyatt, R., Kwong, P. D., Desjardins, E., Sweet, R. W., Robinson, J., Hendrickson, W. A., and Sodroski, J. G. (1998) The antigenic structure of the HIV gp120 envelope glycoprotein, *Nature* **393**, 705-711.
5. Wei, X., Decker, J. M., Wang, S., Hui, H., Kappes, J. C., Wu, X., Salazar-Gonzalez, J. F., Salazar, M. G., Kilby, J. M., Saag, M. S., Komarova, N. L., Nowak, M. A., Hahn, B. H., Kwong, P. D., and Shaw, G. M. (2003) Antibody neutralization and escape by HIV-1, *Nature* **422**, 307-312.
6. Moore, J. P., and Sodroski, J. (1996) Antibody cross-competition analysis of the human immunodeficiency virus type 1 gp120 exterior envelope glycoprotein, *J. Virol.* **70**, 1863-1872.
7. Stewart-Jones, G. B., Soto, C., Lemmin, T., Chuang, G. Y., Druz, A., Kong, R., Thomas, P. V., Wagh, K., Zhou, T., Behrens, A. J., Bylund, T., Choi, C. W., Davison, J. R., Georgiev, I. S., Joyce, M. G., Kwon, Y. D., Pancera, M., Taft, J., Yang, Y., Zhang, B., Shivatare, S. S., Shivatare, V. S., Lee, C. C., Wu, C. Y., Bewley, C. A., Burton, D. R., Koff, W. C., Connors, M., Crispin, M., Baxa, U., Korber, B. T., Wong, C. H., Mascola, J. R., and Kwong, P. D. (2016) Trimeric HIV-1-Env Structures Define Glycan Shields from Clades A, B, and G, *Cell* **165**, 813-826.
8. Burton, D. R., and Mascola, J. R. (2015) Antibody responses to envelope glycoproteins in HIV-1 infection, *Nat Immunol* **16**, 571-576.
9. Crispin, M., and Doores, K. J. (2015) Targeting host-derived glycans on enveloped viruses for antibody-based vaccine design, *Curr. Opin. Virol.* **11**, 63-69.
10. Shingai, M., Donau, O. K., Plishka, R. J., Buckler-White, A., Mascola, J. R., Nabel, G. J., Nason, M. C., Montefiori, D., Moldt, B., Poignard, P., Diskin, R., Bjorkman, P. J., Eckhaus, M. A., Klein, F., Mouquet, H., Cetrulo Lorenzi, J. C., Gazumyan, A., Burton, D. R., Nussenzweig, M. C., Martin, M. A., and Nishimura, Y. (2014) Passive transfer of modest titers of potent and broadly neutralizing anti-HIV monoclonal antibodies block SHIV infection in macaques, *J Exp Med* **211**, 2061-2074.
11. Burton, D. R., Ahmed, R., Barouch, D. H., Butera, S. T., Crotty, S., Godzik, A., Kaufmann, D. E., McElrath, M. J., Nussenzweig, M. C., Pulendran, B., Scanlan, C. N., Schief, W. R., Silvestri, G., Streeck, H., Walker, B. D., Walker, L. M., Ward, A. B., Wilson, I. A., and Wyatt, R. (2012) A Blueprint for HIV Vaccine Discovery, *Cell Host Microbe* **12**, 396-407.
12. Pejchal, R., Doores, K. J., Walker, L. M., Khayat, R., Huang, P. S., Wang, S. K., Stanfield, R. L., Julien, J. P., Ramos, A., Crispin, M., Depetris, R., Katpally, U., Marozsan, A., Cupo, A., Malveste, S., Liu, Y., McBride, R., Ito, Y., Sanders, R. W., Ogohara, C., Paulson, J. C., Feizi, T., Scanlan, C. N., Wong, C. H., Moore, J. P., Olson, W. C., Ward, A. B., Poignard, P., Schief, W. R., Burton, D. R., and Wilson, I. A. (2011) A potent and broad neutralizing antibody recognizes and penetrates the HIV glycan shield, *Science* **334**, 1097-1103.
13. Pancera, M., Shahzad-UI-Hussan, S., Doria-Rose, N. A., McLellan, J. S., Bailor, R. T., Dai, K., Loesgen, S., Louder, M. K., Staube, R. P., Yang, Y., Zhang, B., Parks, R., Eudailey, J., Lloyd, K. E., Blinn, J., Alam, S. M., Haynes, B. F., Amin, M. N., Wang, L. X., Burton, D. R., Koff, W. C., Nabel, G. J., Mascola, J. R., Bewley, C. A., and Kwong, P. D. (2013) Structural basis for diverse N-glycan recognition by HIV-1-neutralizing V1-V2-directed antibody PG16, *Nat Struct Mol Biol* **20**, 804-813.
14. Coss, K. P., Vasiljevic, S., Pritchard, L. K., Krumm, S. A., Glaze, M., Madzorera, S., Moore, P. L., Crispin, M., and Doores, K. J. (2016) HIV-1 glycan density drives the persistence of the mannose patch within an infected individual, *J Virol.*
15. Townsley, S., Mohamed, Z., Guo, W., McKenna, J., Cleveland, B., LaBranche, C., Beaumont, D., Shen, X., Yates, N. L., Pinter, A., Tomaras, G. D., Ferrari, G., Montefiori, D. C., and Hu, S. L. (2016) Induction of heterologous Tier 2 HIV-1 neutralizing and cross-reactive V1/V2-specific antibodies in rabbits by prime-boost immunization, *J Virol.*
16. Crooks, E. T., Tong, T., Chakrabarti, B., Narayan, K., Georgiev, I. S., Menis, S., Huang, X., Kulp, D., Osawa, K., Muranaka, J., Stewart-Jones, G., Destefano, J., O'Dell, S., LaBranche, C., Robinson, J. E., Montefiori, D. C., McKee, K., Du, S. X., Doria-Rose, N., Kwong, P. D., Mascola, J. R., Zhu, P., Schief, W. R., Wyatt, R. T., Whalen, R. G., and Binley, J. M. (2015) Vaccine-Elicited Tier 2 HIV-1 Neutralizing Antibodies Bind to Quaternary Epitopes Involving Glycan-Deficient Patches Proximal to the CD4 Binding Site, *PLoS Pathog* **11**, e1004932.
17. Fournier, J. (2015) A Review of Glycan Analysis Requirements, *BioPharm Int.* **28**, 32-37.

18. Lewis, G. K., DeVico, A. L., and Gallo, R. C. (2014) Antibody persistence and T-cell balance: two key factors confronting HIV vaccine development, *Proc Natl Acad Sci U S A* **111**, 15614-15621.
19. Jardine, J., Julien, J. P., Menis, S., Ota, T., Kalyuzhnyi, O., McGuire, A., Sok, D., Huang, P. S., MacPherson, S., Jones, M., Nieuwsma, T., Mathison, J., Baker, D., Ward, A. B., Burton, D. R., Stamatatos, L., Nemazee, D., Wilson, I. A., and Schief, W. R. (2013) Rational HIV Immunogen Design to Target Specific Germline B Cell Receptors, *Science* **340**, 711-716.
20. de Taeye, S. W., Ozorowski, G., Torrents de la Pena, A., Guttman, M., Julien, J. P., van den Kerkhof, T. L., Burger, J. A., Pritchard, L. K., Pugach, P., Yasmeen, A., Crampton, J., Hu, J., Bontjer, I., Torres, J. L., Arendt, H., DeStefano, J., Koff, W. C., Schuitemaker, H., Eggink, D., Berkhout, B., Dean, H., LaBranche, C., Crotty, S., Crispin, M., Montefiori, D. C., Klasse, P. J., Lee, K. K., Moore, J. P., Wilson, I. A., Ward, A. B., and Sanders, R. W. (2015) Immunogenicity of Stabilized HIV-1 Envelope Trimers with Reduced Exposure of Non-neutralizing Epitopes, *Cell* **163**, 1702-1715.
21. Rudd, P. M., and Dwek, R. A. (1997) Glycosylation: heterogeneity and the 3D structure of proteins, *Crit. Rev. Biochem. Mol. Biol.* **32**, 1-100.
22. Behrens, A. J., Vasiljevic, S., Pritchard, L. K., Harvey, D. J., Andev, R. S., Krumm, S. A., Struwe, W. B., Cupo, A., Kumar, A., Zitzmann, N., Seabright, G. E., Kramer, H. B., Spencer, D. I., Royle, L., Lee, J. H., Klasse, P. J., Burton, D. R., Wilson, I. A., Ward, A. B., Sanders, R. W., Moore, J. P., Doores, K. J., and Crispin, M. (2016) Composition and Antigenic Effects of Individual Glycan Sites of a Trimeric HIV-1 Envelope Glycoprotein, *Cell Rep* **14**, 2695-2706.
23. Go, E. P., Herschhorn, A., Gu, C., Castillo-Mendez, L., Zhang, S., Mao, Y., Chen, H., Ding, H., Wakefield, J. K., Hua, D., Liao, H. X., Kappes, J. C., Sodroski, J., and Desaire, H. (2015) Comparative Analysis of the Glycosylation Profiles of Membrane-Anchored HIV-1 Envelope Glycoprotein Trimers and Soluble gp140, *J Virol* **89**, 8245-8257.
24. Behrens, A. J., Harvey, D. J., Milne, E., Cupo, A., Kumar, A., Zitzmann, N., Struwe, W. B., Moore, J. P., and Crispin, M. (2016) Molecular architecture of the cleavage-dependent mannose patch on a soluble HIV-1 envelope glycoprotein trimer, *J Virol*.
25. Zhu, Z., Go, E. P., and Desaire, H. (2014) Absolute quantitation of glycosylation site occupancy using isotopically labeled standards and LC-MS, *J Am Soc Mass Spectrom* **25**, 1012-1017.
26. Go, E. P., Liao, H. X., Alam, S. M., Hua, D., Haynes, B. F., and Desaire, H. (2013) Characterization of host-cell line specific glycosylation profiles of early transmitted/founder HIV-1 gp120 envelope proteins, *J. Proteome Res.* **12**, 1223-1234.
27. Hernandez, H., and Robinson, C. V. (2007) Determining the stoichiometry and interactions of macromolecular assemblies from mass spectrometry, *Nat Protoc* **2**, 715-726.
28. Elbein, A. D., Tropea, J. E., Mitchell, M., and Kaushal, G. P. (1990) Kifunensine, a potent inhibitor of the glycoprotein processing mannosidase I, *J Biol Chem* **265**, 15599-15605.
29. Gault, J., Donlan, J. A., Liko, I., Hopper, J. T., Gupta, K., Housden, N. G., Struwe, W. B., Marty, M. T., Mize, T., Bechara, C., Zhu, Y., Wu, B., Kleanthous, C., Belov, M., Damoc, E., Makarov, A., and Robinson, C. V. (2016) High-resolution mass spectrometry of small molecules bound to membrane proteins, *Nat. Methods* **13**, 333-336.
30. Yang, Y., Barendregt, A., Kamerling, J. P., and Heck, A. J. (2013) Analyzing protein micro-heterogeneity in chicken ovalbumin by high-resolution native mass spectrometry exposes qualitatively and semi-quantitatively 59 proteoforms, *Anal. Chem.* **85**, 12037-12045.
31. Rosati, S., van den Bremer, E. T., Schuurman, J., Parren, P. W., Kamerling, J. P., and Heck, A. J. (2013) In-depth qualitative and quantitative analysis of composite glycosylation profiles and other micro-heterogeneity on intact monoclonal antibodies by high-resolution native mass spectrometry using a modified Orbitrap, *mAbs* **5**, 917-924.
32. Parsons, T. B., Struwe, W. B., Gault, J., Yamamoto, K., Taylor, T. A., Raj, R., Wals, K., Mohammed, S., Robinson, C. V., Benesch, J. L., and Davis, B. G. (2016) Optimal Synthetic Glycosylation of a Therapeutic Antibody, *Angew Chem Int Ed Engl* **55**, 2361-2367.
33. Chang, V. T., Crispin, M., Aricescu, A. R., Harvey, D. J., Nettleship, J. E., Fennelly, J. A., Yu, C., Boles, K. S., Evans, E. J., Stuart, D. I., Dwek, R. A., Jones, E. Y., Owens, R. J., and Davis, S. J. (2007) Glycoprotein structural genomics: solving the glycosylation problem, *Structure* **15**, 267-273.
34. Crispin, M., Bowden, T. A., Coles, C. H., Harlos, K., Aricescu, A. R., Harvey, D. J., Stuart, D. I., and Jones, E. Y. (2009) Carbohydrate and domain architecture of an immature antibody glycoform exhibiting enhanced effector functions, *J. Mol. Biol.* **387**, 1061-1066.
35. Scanlan, C. N., Ritchie, G. E., Baruah, K., Crispin, M., Harvey, D. J., Singer, B. B., Lucka, L., Wormald, M. R., Wentworth, P., Jr., Zitzmann, N., Rudd, P. M., Burton, D. R., and Dwek, R. A. (2007) Inhibition of mammalian glycan biosynthesis produces non-self antigens for a broadly neutralising, HIV-1 specific antibody, *J. Mol. Biol.* **372**, 16-22.
36. Harvey, D. J., Scarff, C. A., Edgeworth, M., Struwe, W. B., Pagel, K., Thalassinou, K., Crispin, M., and Scrivens, J. (2016) Travelling-wave ion mobility and negative ion fragmentation of high-mannose N-glycans, *J Mass Spectrom* **51**, 219-235.
37. Stephenson, K. E., D' Couto, H. T., and Barouch, D. H. (2016) New concepts in HIV-1 vaccine development, *Curr. Opin. Immunol.* **41**, 39-46.
38. Stansell, E., Panico, M., Canis, K., Pang, P. C., Bouche, L., Binet, D., O'Connor, M. J., Chertova, E., Bess, J., Lifson, J. D., Haslam, S. M., Morris, H. R., Desrosiers, R. C., and Dell, A. (2015) Gp120 on HIV-1 Virions Lacks O-Linked Carbohydrate, *PLoS One* **10**, e0124784.
39. Lingg, N., Zhang, P., Song, Z., and Bardor, M. (2012) The sweet tooth of biopharmaceuticals: importance of recombinant protein glycosylation analysis, *Biotechnol J* **7**, 1462-1472.

HIV Glycan Complexity




i. Kifunensine

ii. Intact MS



Glycan Occupancy



Abstract Graphic

176x67mm (149 x 149 DPI)